

CHILLI FARMING

Farming time

- In the hills chilli is grown in the rainy season.
- In the low hills and valleys it is grown in the winter season.
- Chilli cannot bear extreme hot and extreme cold temperature.


Month/ Region	Apr/ May	May /Jun	Jun/ Jul	Jul/ Aug	Aug/ Sep	Sep/ Oct	Oct/ Nov	Nov/ Dec	Dec/ Jan	Jan/ Feb	Feb/ Mar	Mar/ Apr
High hills	1	**	華	華	華							
Mid hills	華	奎	牽	奎	華	華						華
Valley/ lowlan ds					1111	華	棄	華	棄	棄	華	華

Breed

- Suryamukhi
- Jwala
- Akabarey

Nursery

- In the hills, chilli saplings should be cultivated inside plastic greenhouses.
- In the valleys, raised nursery beds with shed should be used.






Practical Action, The Schumacher Centre, Bourton on Dunsmore, Rugby, Warwickshire, CV23 9QZ, UK T $_{+44}$ (0)1926 634400 | F $_{+44}$ (0)1926 634401 | E infoserv@practicalaction.org.uk | W $_{www.practicalaction.org}$

Seed and Sapling Rate

- Seeds: 60 gm per 0.05 hectare land
- Saplings: 4,000–5,000 plants per 0.05 hectare land
- Good yield can be had from the use of quality seeds.





Land Preparation

- In the winter and spring season, the chilli crop can be cultivated in plain and flat land.
- In the rainy season the crop can be cultivated in ridges.
- Ridges should be built only after deep tilling.





Fertilizer


• Fifty baskets (approximately 1.2 MT) of well-decomposed manure should be used per 0.05 hectare land.








• If available, 2 kg of urea, 4.5 kg of D.A.P. and 3.5 kg of Muriate of potash can be used for every 0.05 hectare land during the final stage of land preparation.



Transplanting of Sapling

- During winter, saplings should be transplanted when they have 5–6 leaves.
- In the rainy season, saplings should be transplanted when they have 7–8 leaves.



Weed Control

Weeds should not be allowed to grow when farming chillis.











Additional Fertilizer

• A mixture of cattle urine and water should be applied at the rate of 1 litre per 1 sq m of land every 15 days.




- In case of use of chemical fertilizers, 4 and 8 days after transplanting the saplings, 1,500 gm of urea per 0.05 hectare land should be applied around the plants at a time.
- After applying the manure, the crop should be covered with soil.
- If the soil lacks moisture, additional fertilizer should be applied and irrigation should be done.



Harvesting

- Chilli should be plucked while they are green and shiny. Their stem should be hard and should easily break while plucking.
- Ripened chilli is usually dried.







Storing

• Usually, chillies are properly dried in the sun and later used as spice.



(Source: Krishak Sahayogi Pustika: Nepalka Pramukh Tarakari tatha Khadanna Bali Kheti Prabidhi)

Written by READ Nepal and Practical Action Nepal and translated by Binaya Parajuli in September 2015.

Note: Any unauthorized use and sale of this publication for commercial purpose is strictly prohibited without the consent of the publisher. All the information provided in this publication has been published in consultation with experts and information published by other organizations. Steps that can be taken by communities on their own effort has been emphasized in this publication. For special conditions, related experts should be consulted.

Practical Action Nepal

P O Box 15135 Narayan Gopal Sadak Paninpokhari, Maharajgunj Kathmandu Nepal

Tel: +00 977 1 4423 639 Fax: +00 977 1 442 3632

E-mail: infoserv@practicalaction.org.uk
Website: http://www.practicalaction.org.uk

Practical Action is a development charity with a difference. We know the simplest ideas can have the most profound, life-changing effect on poor people across the world. For over 40 years, we have been working closely with some of the world's poorest people - using simple technology to fight poverty and transform their lives for the